General Disclaimer

One or more of the Following Statements may affect this Document

- This document has been reproduced from the best copy furnished by the organizational source. It is being released in the interest of making available as much information as possible.
- This document may contain data, which exceeds the sheet parameters. It was furnished in this condition by the organizational source and is the best copy available.
- This document may contain tone-on-tone or color graphs, charts and/or pictures, which have been reproduced in black and white.
- This document is paginated as submitted by the original source.
- Portions of this document are not fully legible due to the historical nature of some
 of the material. However, it is the best reproduction available from the original
 submission.

Produced by the NASA Center for Aerospace Information (CASI)

SC507.71AR
Copy No.

LUNAR SAMPLE ANALYSIS

CR-171 851

ANNUAL REPORT 13 FOR THE PERIOD January 1, 1984 through December 31, 1984

(NASA-CR-171851) LUNAE SAMFIE ANALYSIS
Annual Report, 1 Jan. - 31 Dec. 1984
(Bockwell International Science Center) 4 p
BC A02/MF A01
Unclas
G3/91
14373

CONTRACT NO. NAS9-11539

Prepared for

NASA-Lyndon B. Johnson Space Center Science & Applications Directorate Houston, Texas 77058

> R.M. Housley Principal Investigator

FEBRUARY 1985

Approved for public release; distribution unlimited



7

A Michigan & ... 12

٠,	•	Unclassi	fied			
	SECURITY CLASSIFICATION OF THIS PAGE					

	REPORT DOCUME	NTATION PAGE	<u> </u>		ż	
18 REPORT SECURITY CLASSIFICATION	1b. RESTRICTIVE MARKINGS					
Unclassified	None					
28. SECURITY CLASSIFICATION AUTHORITY	Approved for public release; distribution					
76 DECLASSIFICATION/DOWNGRADING SCHEE	unlimited.					
4 PERFORMING ORGANIZATION REPORT NUM SC507.71AR	5. MONITORING ORGANIZATION REPORT NUMBER(S)					
Rockwell International/ Science Center	78. NAME OF MONITORING ORGANIZATION					
P.O. Box 1085 Thousand Oaks, CA 91360	7b. ADGRESS (City, State and ZIP Code)					
NAME OF FUNDING/SPONSORING ORGANIZATION NASA Lyndon B. Johnson	PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER NAS9-11539					
Bc ADDRESS (City, State and ZIP Code)		10 SOURCE OF FUNDING NOS				
Science & Applications Dire Houston, Texas 77058	PROGRAM ELEMENT NO.	PROJECT NO.	TASK No.	WORK UNIT		
11 TITLE (Include Security Classification) LUNAR SAMPLE ANALYSIS						
12. PERSONAL AUTHORIS! R.M. Housley						
Annual Report No. 13 FROM O	1/01/84to 12/31/8	January 2	RT (Yr. Mo. Day) 7. 1985	15 PAGE	COUNT	
16. SUPPLEMENTARY NOTATION						
17 COSATI CODES	IB. SUBJECT TERMS IC	S (Continue on reverse if necessary and identify by block number)				
FIELD GROUP SUB GR	Allende, ALHA 77003, Chondrule, Alteration Unequilibrated, Calcium, Pyroxene, Ulivine					
Recent results on the anta standing the alteration pr chondrites are presented. microscope results confirm alteration.	rctic meteorite ocesses which pr Also included a	ALHA 77003 wh oduced matrix are additional	in unequil scanning e	lectron	er-	
20 DISTRIBUTION/AVAILABILITY OF ABSTRA	21. ABSTRACT SECURITY CLASSIFICATION					
UNCLASSIFIED/UNLIMITED - BAME AS RPT.	Unclass	ified				
22a. NAME OF RESPONSIBLE INDIVIDUAL R.M. Housley	22b TELEPHONE N (Include Area Co 805/373-42	del	22c, OFFICE SY	ИВО L		

DD FORM 1473, 83 APR

EDITION OF 1 JAN 73 IS OBSOLETE.

Unclassified

ADDITIONAL EVIDENCE POINTING TO THE FORM TION OF ALLENDE MATRIX BY IN SITU ALTERATION. R. M. Housley, Rockwell International Science Center, Thousand Oaks, CA 91360.

"One of the important, and yet frequently violated ground rules of this game is that all facts, all interpretations, and especially all premises be critically reviewed from time to time." Anders, 1964, (2).

It has probably always been evident (1) that it is much easier to produce material resembling matrix in unequilibrated chondrites by parent body alter-

ation processes than it is in any space environment.

The basic difficulty in producing matrix in the solar nebula comes from the long time it would take to form highly oxidized and hydrated silicates at the temperatures where they would become thermodynamically stable. This would require a long cool quiescent interval between the condensation of most of the material and the accreation of planetary bodies. It has also been known for more than 20 years (2,3) that this difficulty can easily be avoided by forming the matrix in parent bodies, where isolation from hydrogen permits oxidation at higher temperatures.

Despite all this it has seemed convenient until recently to regard the bulk of the matrix ir unequilibrated meteorites as a primary constituent, since this appeared to provide a natural explanation for the systematic depletions in moderately volalite elements such as Zn which are observed when the

other meteorite classes are compared to CI chondrites (2,4).

Serious difficulties with this hypothesis become evident though when one carefully considers the analyses of carbonaceous chondrite matrices reported by McSween and Richardson (5). They found substantial differences in matrix compositions from 100 μm spot to 100 μm spot within the same meteorite as well as in the average matrix compositions between different meteorites of the same class. Similar results for the ordinary chondrites were later reported by Huss et. al. (6). Analogous striking differences in matrix grain size and texture are also readily apparent.

These observations seem incompatible with the hypothesis of a single primative volatile rich matrix material. Each meteorite even within a given class would appear to require its own special matrix which would then have to be mixed in proper proportions with the other constituents to achieve the overall composition appropriate to the class. Such complications are completely

circumvented if matrix resulted form parent body alteration.

We have previously reported (7) extensive scanning electron microscope evidence showing that much matrix olivine in Allende was produced by a reaction

equivalent to MgSiO₃ + Fe + $\frac{1}{2}$ O₂ \rightarrow MgFeSi₂O₄.


Following up a clue mentioned by Peck (8) we now present in Fig. 1 new evidence showing that much additional matrix olivine, as well as, Fe-rich clinopyroxene was produced by in situ alteration of a pre-existing Ca-rich mineral. The backscattered electron images of typical matrix regions compared to the Ca elemental maps of the corresponding regions seem to allow no other simple interpretation. The most probable Ca-rich precursor appears to be diopside. We have recently reported a strong tendency for diopside in the less altered meteorite ALHA 77003 to show alteration to Fe-rich olivine (9).

Besides providing additional evidence for matrix formation by alteration in Allende the current results provide strong evidence against post alteration regolith processes or impact mixing. They seem incompatible with any scenario

which requires the formation of matrix as small grains in space (10). (1) Urey (1961) JGR $\underline{66}$, 1988; (2) Anders (1964) Space Science Reviews 3, 583; (3) Wood (1963) Icarus $\underline{2}$, 152; (4) Wasson and Chou (1974) Meteoritics $\underline{9}$, 69;

R. M. Housley

(5) McSween and Richardson (1977) GCA 41, 1145; (6) Huss et. al. (1981) GCA 45, 33; (7) Housley and Cirlin (1983) Chondrules and their Origins. 145; (8) Peck (1983) Meteoritics 18, 373; (9) Housley (1984) Meteoritics 19, in press; (10) Kornacki and Wood (1984) GCA 48, 1963.



ORIGINAL TITL OF POOR QUALITY